

Lecture 9: Polynomials

Why Only Have One Nomial?

1/21

What Is a Polynomial?

High school: $p(x) = c_d x^d + c_{d-1} x^{d-1} + \dots + c_1 x + c_0$

- ▶ $d \in \mathbb{N}$ is the *degree*
- ▶ c_d, \dots, c_0 are the *coefficients*

This is *coefficient representation*

Need $d + 1$ coefficients to define deg d polynomial

Today: see *value representation*

Need $d + 1$ function values to define deg d poly

Today, prove that these are equivalent!

2/21

Polynomial Long Division

Theorem: Let $p(x)$, $d(x)$ be polys. Then $\exists q(x)$, $r(x)$ st $p(x) = q(x)d(x) + r(x)$ and $\deg(r) < \deg(p)$.

Same idea as elementary school long division!

$$\begin{array}{r} x^2 x^2 + 3x x^2 + 3x - 1 \\ x^2 - 1 \overline{) x^4 + 3x^3 - 2x^2 + 0x + 4} \\ \underline{-(x^4 + 0x^3 - x^2)} \\ 3x^3 - x^2 + 0x \\ \underline{-(3x^3 + 0x^2 - 3x)} \\ -x^2 + 3x + 4 \\ \underline{-(-x^2 + 0x + 1)} \\ 3x + 3 \end{array}$$

3/21

Factoring Roots

Lemma: Suppose $p(a) = 0$. Then can write $p(x) = (x - a)q(x)$ st $\deg(q) = \deg(p) - 1$.

Proof:

- ▶ Divide $p(x)$ by $(x - a)$ as before
- ▶ $p(x) = (x - a)q(x) + r(x)$
- ▶ $0 = p(a) = (a - a)q(a) + r(a) = r(a)$
- ▶ $\deg(r) < \deg(x - a) = 1$, so $r(x)$ a constant
- ▶ Only possibility: $r(x) = 0!$
- ▶ Thus $p(x) = (x - a)q(x)$

4/21

Number of Groots

Theorem: Non-zero deg d poly has $\leq d$ roots.

Proof:

- ▶ By induction on d .
- ▶ Base Case ($d = 0$): constant poly, no roots
- ▶ Suppose true for degree k
- ▶ Let $p(x)$ have degree $k + 1$
- ▶ If p has no roots, done
- ▶ Else can factor as $(x - a)q(x)$
- ▶ 1 root from $(x - a)$, $\leq k$ from $q(x)$
- ▶ Total $\leq k + 1$ roots

5/21

Limited Agreement

Theorem: Distinct deg d polys agree on $\leq d$ points

Proof:

- ▶ Let $p(x)$ and $q(x)$ be distinct, $\deg \leq d$
- ▶ $p(x) = q(x)$ iff $p(x) - q(x) = 0$
- ▶ Note: $p - q$ is non-zero, $\deg \leq d$
- ▶ So $p - q$ has $\leq d$ roots
- ▶ Means $\leq d$ values of x st $p(x) = q(x)!$

Means $d + 1$ values enough to define polynomial

But do any $d + 1$ points work?

6/21

First Interpolation

Want degree 1 poly through (4, 2) and (7, 0)

Recall: slope = rise/run

Here: slope = $(0 - 2)/(7 - 4) = -\frac{2}{3}$

So $p(x) = -\frac{2}{3}x + c$

Choose c st $p(4) = -\frac{2}{3} \cdot 4 + c = 2$

So $c = 2 + \frac{8}{3} = \frac{14}{3}$

So $-\frac{2}{3}x + \frac{14}{3}$ is unique degree 1 poly!

7 / 21

Bigger Interpolation

Want degree 2 through (0, -1), (2, 9), (-1, -3)

Rise/run trick only works for degree 1...



$$\Delta_0(x) = -\frac{1}{2}x^2 + \frac{1}{2}x + 1$$

$$\Delta_2(x) = \frac{1}{6}x^2 + \frac{1}{6}x$$

$$\Delta_{-1}(x) = \frac{1}{3}x^2 - \frac{2}{3}x$$

Take $p(x) = -1\Delta_0(x) + 9\Delta_2(x) - 3\Delta_{-1}(x)$

Works out to $(\frac{1}{2} + \frac{3}{2} - 1)x^2 + (-\frac{1}{2} + \frac{3}{2} + 2)x - 1$

So $p(x) = x^2 + 3x - 1$

8 / 21

Finidng Δ

Goal: $\Delta_0(x)$ st $\Delta_0(0) = 1, \Delta_0(2) = \Delta_0(-1) = 0$

Last two easy: take $q_0(x) = (x - 2)(x + 1)$

Note: $q_0(0) = (0 - 2)(0 + 1) = -2$

Take $\Delta_0(x) = -\frac{1}{2}q_0(x)$

Gives $\Delta_0(x) = -\frac{1}{2}(x^2 - x - 2) = -\frac{1}{2}x^2 + \frac{1}{2}x + 1$

For 2, take $q_2(x) = (x - 0)(x + 1) = x^2 + x$

$\Delta_2(x) = \frac{q_2(x)}{q_2(2)} = \frac{x^2 + x}{6} = \frac{1}{6}x^2 + \frac{1}{6}x$

For -1, take $q_{-1} = (x - 0)(x - 2) = x^2 - 2x$

$\Delta_{-1}(x) = \frac{q_{-1}(x)}{q_{-1}(-1)} = \frac{x^2 - 2x}{3} = \frac{1}{3}x^2 - \frac{2}{3}x$

9 / 21

Lagrange Interpolation

Theorem: Given points $(x_1, y_1), \dots, (x_{d+1}, y_{d+1})$, can construct deg (at most) d poly through them.

Proof:

- ▶ Suppose have polys $\Delta_i(x)$ st
 - ▶ $\Delta_i(x_i) = 1$
 - ▶ $\Delta_i(x_j) = 0$ for $j \neq i$
- ▶ Take $p(x) = y_1\Delta_1(x) + \dots + y_{d+1}\Delta_{d+1}(x)$
- ▶ To construct $\Delta_i(x)$:
 - ▶ Take $q_i(x) = \prod_{j \neq i} (x - x_j)$
 - ▶ Let $\Delta_i(x) = \frac{q_i(x)}{q_i(x_i)}$

Note similarities to CRT!

10 / 21

Lagrange Example

Find deg 2 poly through (1, 6), (6, 1), (7, 0)

$$\Delta_1(x) = \frac{(x-6)(x-7)}{(1-6)(1-7)} = \frac{1}{30}x^2 - \frac{13}{30}x + \frac{42}{30}$$

$$\Delta_6(x) = \frac{(x-1)(x-7)}{(6-1)(6-7)} = -\frac{1}{5}x^2 + \frac{8}{5}x - \frac{7}{5}$$

$$\Delta_7(x) = \frac{(x-1)(x-6)}{(7-1)(7-6)} = \frac{1}{6}x^2 - \frac{7}{6}x + 1$$

So take $p(x) = 6\Delta_1(x) + 1\Delta_6(x) + 0\Delta_7(x)$

$$6\Delta_1(x) = \frac{1}{5}x^2 - \frac{13}{5}x + \frac{42}{5}$$

$$p(x) = (\frac{1}{5}x^2 - \frac{13}{5}x + \frac{42}{5}) + (-\frac{1}{5}x^2 + \frac{8}{5}x - \frac{7}{5}) = -x + 7$$

Notice: doesn't have to be degree *exactly* 2!

11 / 21

Break

Break time! Talk to your neighbors!

Today's Discussion Question:

What is your favorite breakfast food?

12 / 21

Get Real

So far, working with polynomials in \mathbb{R}

Calculations tend to get messy

Issues with finite precision on computers!

What properties of \mathbb{R} did we actually use?

- ▶ Ability to add, multiply, subtract
- ▶ Division by non-zero numbers
- ▶ Product of non-zero numbers is non-zero

These properties hold in any *field*

Numbers modulo a prime is a field!

13 / 21

Finite Fields

Numbers mod p often denoted $GF(p)$ ¹

Note: is important that p is a prime!

Ex: Consider $(x-2)(x-3)$ modulo 6

- ▶ Degree two polynomial
- ▶ But four roots: 0, 2, 3, 5!

Ex: No deg 1 poly through (0,0) and (3,1) mod 6

- ▶ Go through (0,0) means $c_0 = 0$
- ▶ Go through (3,1) means $c_1 \cdot 3 \equiv 1 \pmod{6}$

¹"GF" stands for "Galois Field"

14 / 21

Finite Field Lagrange

Want deg 2 poly mod 7 through (0,3), (2,2), (3,0)

$$q_0(x) = (x-2)(x-3) = x^2 - 5x + 6 \equiv x^2 + 2x + 6$$

$$q_0(0) = 6, \text{ so } q_0(0)^{-1} \equiv 6 \pmod{7}$$

$$\Delta_0(x) \equiv 6(x^2 + 2x + 6) \equiv 6x^2 + 5x + 1 \pmod{7}$$

$$q_2(x) = (x-0)(x-3) = x^2 - 3x \equiv x^2 + 4x \pmod{7}$$

$$q_2(2) = -2 \equiv 5 \pmod{7}, \text{ so } q_2(2)^{-1} \equiv 3 \pmod{7}$$

$$\Delta_2(x) \equiv 3(x^2 + 4x) \equiv 3x^2 + 5x \pmod{7}$$

Don't have to calculate $\Delta_3(x)$ — multiplied by zero!

$$\begin{aligned} \text{Take } p(x) &= 3\Delta_0(x) + 2\Delta_2(x) + 0\Delta_3(x) \\ &\equiv (4x^2 + x + 3) + (6x^2 + 3x) \equiv 3x^2 + 4x + 3 \pmod{7} \end{aligned}$$

15 / 21

Counting Polynomials

Suppose I know $p(1) = 5$ and $p(2) = 3$.
How many deg ≤ 2 polynomials could p be?

Polynomial fully defined by 3rd point
Equiv: how many possible values for $p(0)$?

In \mathbb{R} , infinitely many...not too interesting

In $GF(q)$, q possibilities!

16 / 21

Shhh, It's a Secret

The password to my computer is 1234.

If something bad happens, want staff to unlock it
But dangerous to just give out my password

Idea: if k of n staff members agree, can unlock
If fewer than k , unable to

Shamir's Secret Sharing Scheme:

- ▶ Choose random deg $k-1$ poly st $p(0) = 1234$
 - ▶ Can choose points and interpolate
 - ▶ Or can choose coefficients
- ▶ Distribute $p(i)$ to i th staff member ($1 \leq i \leq n$)

17 / 21

Shhh-amir Properties

Claim: k staff members can recover password

Proof:

- ▶ Any k points on p fully define polynomial
- ▶ Use Lagrange to interpolate; evaluate $p(0)$

Claim: Only $k-1$ staff members get nothing

Proof:

- ▶ Have $k-1$ known points
- ▶ Any value of $p(0)$ gives potential polynomial
- ▶ All values consistent with known points!

18 / 21

Secret Sharing Example

Suppose my secret is 4.

Want to make sure any 3 of the 12 TAs can find it.

What prime should I work modulo?

Eventually give out $p(1)$, $p(2)$, ..., $p(12)$

To ensure distinct, need prime 13 or larger!

(Also need prime bigger than secret)

Choose polynomial $x^2 + 4 \pmod{13}$

Give out $p(1) = 5$, $p(2) = 8$, ..., $p(12) = 5$

Exercise: choose 3 pts, check Lagrange gives $x^2 + 4$

19 / 21

Hierarchical Secret Sharing

Can modify protocol for more complicated setups

Ex: Need Elizabeth + k TAs to unlock

Idea: have nested secret sharing

- ▶ Password is root of degree 1 poly $p(x)$
- ▶ $p(1)$ given to Elizabeth
- ▶ $p(2)$ is secret shared by TAs!
- ▶ Give TAs points on $q(x)$ st $q(0) = p(2)$

See more examples of this in discussion

20 / 21

Fin

Next time: error correcting codes!

21 / 21